

# UC Davis

## UC Davis Previously Published Works

### Title

Adverse metabolic effects of dietary fructose: results from the recent epidemiological, clinical, and mechanistic studies.

### Permalink

<https://escholarship.org/uc/item/2j3108km>

### Journal

Current opinion in lipidology, 24(3)

### ISSN

0957-9672

### Authors

Stanhope, Kimber L  
Schwarz, Jean-Marc  
Havel, Peter J

### Publication Date

2013-06-01

### DOI

10.1097/mol.0b013e3283613bca

Peer reviewed

Published in final edited form as:

*Curr Opin Lipidol.* 2013 June ; 24(3): 198–206. doi:10.1097/MOL.0b013e3283613bca.

## Adverse metabolic effects of dietary fructose: Results from recent epidemiological, clinical, and mechanistic studies

**Kimber L. Stanhope,**

Department of Molecular Biosciences (K.L.S., P.J.H.), School of Veterinary Medicine, and Department of Nutrition (K.L.S., P.J.H.), University of California, Davis (UCD), Davis, California, 95616

**Jean-Marc Schwarz,** and

Department of Basic Sciences (J.M.S.), College of Osteopathic Medicine, Touro University, Vallejo, California 94592; and Department of Medicine (J.M.S.), University of California San Francisco, San Francisco, California, 94143

**Peter J. Havel**

Department of Molecular Biosciences (K.L.S., P.J.H.), School of Veterinary Medicine, and Department of Nutrition (K.L.S., P.J.H.), University of California, Davis (UCD), Davis, California, 95616

### Abstract

**Purpose of review—**The effects of dietary sugar on risk factors and processes associated with metabolic disease remains a controversial topic, with recent reviews of the available evidence arriving at widely discrepant conclusions.

**Recent findings—**There are many recently published epidemiological studies that provide evidence that sugar consumption is associated with metabolic disease. Three recent clinical studies, which investigated the effects of consuming relevant doses of sucrose or high fructose corn syrup along with *ad libitum* diets, provide evidence that consumption of these sugars increase risk factors for cardiovascular disease (CVD) and metabolic syndrome. Mechanistic studies suggest that these effects result from the rapid hepatic metabolism of fructose catalyzed by fructokinase C, which generates substrate for *de novo* lipogenesis and leads to increased uric acid levels. Recent clinical studies investigating the effects of consuming less sugar, via educational interventions or by substitution of sugar-sweetened beverages for non-calorically sweetened beverages, provide evidence that such strategies have beneficial effects on risk factors for metabolic disease or on BMI in children.

**Summary—**The accumulating epidemiological evidence, direct clinical evidence, and the evidence suggesting plausible mechanisms support a role for sugar in the epidemics of metabolic syndrome, CVD and type 2 diabetes.

## Keywords

Fructose; sucrose; high fructose corn syrup; sugar; metabolic disease

## Introduction

In August 2009, the American Heart Association Nutrition Committee recommended that women consume no more than 100 kcal/day and men consume no more than 150 kcal/day of added sugar (1). In June 2010, the Report of the Dietary Guidelines Advisory Committee (DGAC) on the Dietary Guidelines for Americans 2010, suggested a maximal intake level of 25% or less of total energy from added sugars (2). While this latter guideline is not recommending people consume 25% of their energy as added sugar, it does imply that after due consideration of the evidence DGAC concluded that consumption of added sugar at this level is not associated with any adverse metabolic effects. The difference in these 2 guidelines, equivalent to almost three 12-ounce servings of soda for the average woman and more than 3.5 for the average man, illustrates the state of controversy that existed concerning the effects of sugar consumption on the development of metabolic disease in 2010.

Are we any nearer a consensus now, 3 years later? The discrepant conclusions summarized below from review articles evaluating the available data to 2012 suggest we are not.

- Data from prospective and intervention studies clearly point to high fructose consumption, mainly in the form of sugar-sweetened beverages (SSB) as risk factor for metabolic diseases in humans (3).
- Although some studies hint towards some potential adverse effects of excessive fructose consumption especially when combined with excess energy intake, the results from clinical trials do not support a significant detrimental effect of fructose on metabolic health when consumed as part of a weight-maintaining diet in amounts consistent with the average-estimated fructose consumption in Western countries (4).
- Intake of free sugar or SSB is a determinant of body weight (5).
- Randomized controlled trials at levels even exceeding normal human consumption have been inconclusive related to SSB and obesity (6).

This review will present the recent epidemiological, clinical, and mechanistic studies pertaining to the effects of dietary sugar on risk factors and processes associated with metabolic disease, and provide the perspective of researchers directly involved in clinical investigations of this topic.

## Epidemiological studies

Recent studies add to the already considerable epidemiological evidence that sugar consumption is associated with metabolic disease.

- Men from the Health Professionals Follow-Up Study in the top quartile of SSB intake had a 20% higher relative risk of coronary heart disease than those in the bottom quartile (7).
- In Hispanic adults, plasma TG, metabolic syndrome and waist circumference were associated with consumption of instant SSB and/or regular soda (8).
- In the Nurses' Health Study II, one serving/d of SSB was associated with increased risk for type 2 diabetes (T2DM) (9).
- In participants from the Nurses' Health Study, Nurses' Health Study II and the Health Professionals Follow-Up Study, SSB consumption was associated with a higher risk of T2DM; coffee intake was associated with a lower risk, irrespective of the caffeine content. (10)
- In an analysis across 94 countries, every additional percentage point of calories from sugar/sweeteners were associated with 5% higher prevalence of diabetes (11).
- An increased consumption of 100-mL/day of SSB was associated with increased HOMA IR and systolic blood pressure among children 85 percentile for BMI in the Quebec Adiposity and Lifestyle Investigation in Youth Study, and with increased systolic blood pressure and waist circumference in children with impaired glucose tolerance (12).
- In healthy adults in Scotland, uric acid levels were positively associated with SSB consumption (13).
- SSB consumption was positively associated with serum uric acid concentrations in adolescents in Taiwan, as was BMI, body fat, and systolic blood pressure. 25% of the 2727 subjects consumed more than 500 ml of SSB/day (14).
- Among participants of the Nurses' Health Study, Nurses' Health Study II and Health Professionals Follow-Up Study substitution of water, coffee, tea, diet beverages, or low-fat milk for one serving of SSB was associated with weight loss, with the greatest effect occurring with water (15). The genetic association with BMI was stronger among participants with higher intake of SSB than those with lower intake (16).
- In adults, frequency of SSB, but not diet beverage, intake was positively associated with proportion of visceral (VAT) to subcutaneous abdominal adipose tissue (SAT) (17).
- In teenagers, fructose intake was associated with VAT, but not SAT (18).

In contrast to the above reports, an analysis of the NHANES 1999-2006 indicated that fructose and non-fructose sugar consumptions at levels representative of the American diet were not associated with indicators of the metabolic syndrome (19).

## Clinical studies – Interventions with increased sugar intake

Recent clinical studies have investigated the effects of fructose or sugar consumption by providing subjects with SSB or control beverages that were consumed along with *ad libitum*

quantities of the subjects' usual diets. The longest of these studies was conducted in Denmark where subjects consumed 1 liter/d of sucrose-sweetened cola (~20% energy requirements (ER)), isocaloric amounts of low-fat milk, 1 liter/d aspartame-sweetened beverages, or 1 liter/d water for 6 months. Body weight at the end of the intervention period was not significantly different from baseline in any group. Subjects consuming sucrose exhibited increased VAT, liver and muscle TG, and fasting TG and cholesterol levels, while the other 3 groups did not. The sucrose-induced increase of liver TG was significantly larger compared with all 3 of the other groups, and the increase in visceral fat was significantly greater compared with the subjects who consumed low-fat milk and who exhibited a comparable change in body weight (Sucrose: +1.3%; Milk: +1.4%) (20).

Aeberli and colleagues have published two recent reports in which young men participating in 6-arm (21) or 4-arm crossover trials (22) consumed low (40 g/d, ~6.5% ER) or moderate (80 g/d, ~13% ER) amounts of fructose, glucose, or sucrose as beverages along with *ad libitum* diets for 3 weeks. In the 6-arm crossover, LDL particle size was reduced compared with baseline during consumption of moderate fructose or sucrose, and redistribution to a more atherogenic LDL subclass distribution was observed after these interventions plus the low fructose intervention. These 3 interventions also increased waist:hip ratio, even though low glucose was the only intervention that resulted in significant weight gain compared with baseline (21). In the 4-arm crossover, total and LDL cholesterol were increased after moderate fructose or sucrose compared with moderate glucose consumption. Hepatic insulin sensitivity, indexed by endogenous glucose production during euglycemic-hyperinsulinemic clamps, was decreased during moderate fructose compared with moderate glucose consumption, while whole body insulin sensitivity was not different (22).

Our group demonstrated that young, healthy subjects consuming 25% ER as HFCS-sweetened beverages for 2 weeks exhibited significant increases of fasting LDL, non-HDL-cholesterol and apolipoprotein B (apoB), and postprandial TG, remnant-cholesterol and -TG, and small dense LDL (sdLDL), which were comparable to those observed in subjects consuming fructose, and greater than those in subjects consuming glucose (23). In contrast, Silbernagel et al. reported that when subjects consumed 150 g/d of fructose or glucose, the only significant difference between groups was an increase of fasting TG concentrations in the fructose group (24). They did not measure postprandial TG, fasting apoB or sdLDL. Given the similarities between this study (24) and our group's study (23) (both parallel arm, subjects of comparable age and BMI, similar amounts of sugar consumed), it is interesting to consider why there were numerous differential effects between glucose and fructose on lipids in our study, and only one in this study. A potential reason for the lack of group differences in the study by Silbernagel et al. was the highly significant weight gain that occurred in subjects consuming glucose ( $+1.7 \pm 0.4$  kg,  $P = 0.001$  vs baseline,  $P = 0.056$  vs fructose), but not in subjects consuming fructose ( $+0.2 \pm 0.6$  kg) (24). This difference may possibly be due to fructose malabsorption (25), which they did not assess. As shown in Figure 1, sub-division of the 48 subjects who participated in our study (23) into groups that gained or did not gain body weight, illustrates that weight gain has a marked effect on the sugar-induced increases of fasting cholesterol, LDL and apoB concentrations.

The studies by Maersk et al. (20), Aeberli et al. (21, 22) and Stanhope et al. (23) provide direct evidence that consumption of sugar can increase risk factors for metabolic disease. These results are relevant to public health in that the sugars investigated included the commonly-consumed sugars (sucrose and HFCS as opposed to pure fructose) and in quantities comparable to that consumed by a significant number of people (21, 22), and within the maximal intake level of 25% suggested by DGAC (2). Obtaining stronger evidence will require clinical trials in which all study food is formulated and provided throughout the investigation to ensure that there are no diet variations between the experimental groups or interventions that may confound results.

## Clinical studies providing mechanistic insights

Other recent clinical studies investigating the effects of hyper-energetic feeding protocols and/or pure fructose have provided mechanistic insights. In an overfeeding study in which diets were supplemented with 1000 kcal/d as candy/SSB, within 3 weeks subjects exhibited increased body weight (+2%), liver fat (+27%) and increased DNL. The increase of DNL was proportional to the increase in liver fat (26). While the lack of a control group prevents differentiating the effects of sugar from those of overfeeding, these results suggest that DNL is involved in the process by which surplus sugar in the context of positive energy balance increases liver fat. Importantly, our group has previously reported that DNL was increased in subjects consuming fructose with energy-balanced, steady state meals, but not in subjects consuming glucose (27).

However, a recent review of fructose metabolism and isotopic tracer studies concluded that a small percentage of ingested fructose (<1%) appears to be directly converted to plasma TG (28). This figure is clearly an underestimation, based on acute feeding studies that do not take into account that DNL-derived lipid can spend from 24 to over 72 hours in the liver prior to being packaged into very low density lipoprotein (VLDL) and secreted into the circulation (29, 30). The actual percentage of fructose converted to fat is difficult to quantify, especially under physiologically relevant meal-fed conditions, and has yet to be determined. Accurate estimations will require assessments of DNL and VLDL production, secretion and clearance using non-steady state tracer kinetic models. However, the above studies and others demonstrate that DNL is upregulated concurrently with fructose-induced postprandial hypertriglyceridemia (27, 31, 32) and liver fat accumulation (26). The seminal study from Donnelly et al. (29) shows that in patients with NAFLD, 26% of both intra-hepatic fat and VLDLTG are made *de novo* (29). Furthermore, when hepatic DNL is induced, not only are new lipids synthesized and non-esterified fatty acids re-esterified, but hepatic lipid oxidation is down-regulated. Our group has recently reported that the subjects who exhibited increased DNL during fructose consumption, also exhibited inhibition of post-meal lipid oxidation (33). These combined events create an imbalance between hepatic lipid “input” and “export”, leading to net intrahepatic fat accumulation. While we did not measure hepatic lipid in these subjects, the 17% decrease in insulin sensitivity (27) supports the concept that hepatic DNL is a mechanism leading to increased hepatic lipid production, hepatic lipid accumulation, and thereby to hepatic insulin resistance (34).

Bortolotti et al. investigated the hypothesis that high dietary protein content would reverse the inhibition of lipid oxidation and the increase in postprandial TG levels caused by fructose consumption. The concurrent feeding of fructose and protein did not increase lipid oxidation, and in opposition to the hypothesis, it increased postprandial TG levels. The authors suggest that the supplemental protein may have enhanced hepatic VLDL synthesis, assembly and secretion (35).

Our group has also recently reported that 10-weeks consumption of fructose, but not glucose, led to significantly increased 24-h uric acid profiles (36), which suggests that the epidemiological associations (13, 14, 37-39) between sugar consumption and uric acid levels are causal. Fasting concentrations of markers of inflammation; monocyte chemoattractant protein-1, plasminogen activator inhibitor-1, and E-selectin; as well as retinol binding protein-4 and the liver enzyme, gamma glutamyl transferase, were also increased in these same subjects consuming fructose (36, 40).

Our recent report of the postprandial glucose and insulin responses in the subjects consuming glucose or fructose for 10 weeks (41) has clinically relevant implications to the potential role of the glycemic index (GI) in metabolic disease risk. The adverse metabolic effects of dietary sugars have been attributed by some to GI (42, 43). The GI of fructose is 23 compared with 100 for glucose. The calculated relative GI of the baseline high complex carbohydrate diet, the high glucose and the high fructose intervention diets, consumed during the 24-h blood collections in our studies, were 64, 83 and 38, respectively. As expected, for this study (27), and our more recent 2-week study (23), the glucose and insulin excursions of the diets paralleled the GI, with exposure being highest on the glucose diet, intermediate on the complex carbohydrate baseline diet, and lowest on the fructose diet. However, it was subjects consuming the high fructose diets with the lowest GI and glycemic exposure, who exhibited increased VAT and decreased insulin sensitivity (27) and increases of LDL, apoB, and postprandial TG (23, 27). In contrast, when subjects consumed high glucose diets, postprandial plasma glucose and insulin excursions increased substantially (23, 27), however insulin sensitivity (27) and postprandial TG exposure, LDL, and apoB remained unchanged (23, 27). Thus, these results do not support the hypothesis that elevated postprandial glucose and/or insulin excursions contribute to dyslipidemia and insulin resistance. They also demonstrate that studies investigating the relationship of dietary carbohydrates to risk factors for metabolic diseases should accurately determine the glucose and fructose contents of the diets. Dietary fructose may be an important contributor to the inconsistent reported effects of dietary GI on metabolic disease risk. It is likely that other differences between high and low GI diets, with alterations in dietary fiber content being the most likely confounder, underlie these inconsistencies. Nonetheless, the available evidence indicates that it is the fructose and not the glucose component of sucrose and HFCS that is primarily responsible for their adverse metabolic effects.

As the prevalence of pediatric obesity and metabolic syndrome increase, investigations of the effects of sugar consumption in children are needed. The 24-h TG profile was measured in children, with or without nonalcoholic fatty liver disease (NAFLD), during consumption of fructose and glucose in crossover feeding trials (44). As previously shown in adults (45, 46), postprandial TG levels were higher during fructose compared with glucose



consumption in all children. Importantly, the fructose-induced increases in TG were higher in children with NAFLD than those without NAFLD (44).

## Clinical studies – interventions to reduce sugar intake

Several recent clinical studies have demonstrated beneficial effects on metabolic parameters by providing subjects with educational programs aimed at reducing sugar/fructose consumption. Obese African-American and Latino adolescents participated in a 16-week nutrition education intervention focused on decreasing added sugar intake to 10% of daily calories and increasing fiber intake. Despite unchanged BMI, subjects exhibited improved insulin sensitivity compared with the control group (47). Goran et al. have recently reviewed the inter-relationship between genetic factors, liver fat, and sugar consumption, (48) which appear to make Latino children (49) and adults (50) particularly vulnerable to the adverse effects of sugar.

In another study, patients with chronic kidney disease followed dietary instructions and lowered fructose consumption to 12 g/d. After 6 weeks, they exhibited significant decreases in fasting insulin, high sensitivity C-reactive protein and soluble intercellular adhesion molecule, and nonsignificant ( $P<0.1$ ) decreases in blood pressure and uric acid levels (51).

Three recent studies provide evidence that dietary education programs designed to reduce sugar and fructose consumption (52), or blinded (53) or unblinded (54) replacement of SSB with noncaloric-sweetened beverages have beneficial effects on BMI in children. Previous studies suggest that consumption of non-caloric sweeteners compared with sucrose or HFCS also has beneficial effects on BMI in adults (55-57). The explanation for these results could be as simple as people tend to over-eat sugar because they like the sweet taste. However, recent studies on the central effects of sugars in the brain (58-60), made possible by functional magnetic resonance imaging technology, suggest the answer could be more complicated. The most recent of these studies reports that consumption of a fructose-sweetened beverage resulted in greater hypothalamic activation, which would be associated with less appetite suppression, than consumption of a glucose-sweetened beverage in young healthy adults (59). Corroborating these results, the subjects recorded significantly higher ratings of fullness and satiety after the glucose, but not fructose, drink (59).

These data provide a plausible link for the associations between the consumption of sugar-sweetened beverages and body weight gain (16, 61-63). Confirming this link will be important not only for combating the obesity epidemic, but also the obesity-related increases of metabolic syndrome, CVD and T2DM. Recent research (20-23, 27) and older research (64-69) demonstrate that consumption of fructose and fructose-containing sugars has adverse effects on risk factors for metabolic disease that are independent of body weight gain. These results, the results in Figure 1, and the potential link between sugar consumption and body weight gain suggest that sugar promotes the development of metabolic disease through two mechanisms; directly via the adverse effects of fructose on lipid and carbohydrate metabolism, and indirectly by promoting body weight gain (See Figure 2).



## Animal and *in vitro* studies investigating mechanisms of the metabolic effects of fructose

Numerous recent studies, more than can be discussed in this review, have investigated the links between dietary fructose and adverse metabolic effects in animal models or *in vitro* systems. We have included a few studies that provide insight into the underlying mechanisms by which fructose induces lipid dysregulation.

The initial phosphorylation of dietary fructose is largely catalyzed by fructokinase, which is not regulated by hepatic energy status. This results in high levels of fructose uptake by the liver with little of the ingested fructose reaching the systemic circulation. Ishimoto et al. have provided compelling evidence concerning the importance of fructokinase in mediating the adverse effects of fructose (70). They report that fructokinase exists as two isoforms: fructokinase A which is widely distributed and has low affinity for fructose; and fructokinase C, which is expressed primarily in liver, intestine, and kidney and has high affinity for fructose. They demonstrated that fructose-induced metabolic syndrome is prevented in knockout mice lacking both isoforms, but is exacerbated in fructokinase A knockout mice compared with wild-type mice (70). These results demonstrate that fructokinase C is the key driver of the adverse effects induced by fructose. It also suggests that fructokinase A offers some protection against the adverse effects of fructose by allowing for some fructose metabolism in peripheral tissues.

Recent studies have investigated the regulation and effects of fructose-induced DNL. Erion et al. compared the effect of treatment with carbohydrate response element-binding protein (ChREBP) and control antisense oligonucleotides (ASO) in rats fed high fructose or high fat diets (71). Treatment with ChREBP ASO decreased plasma TG concentrations compared with control ASO in both diet groups, but hepatic lipid content and insulin sensitivity were unaffected. The reduction in plasma TG was more pronounced in the fructose-fed group and attributed to measured decreases of hepatic expression of fructokinase, lipogenic genes, and microsomal TG transfer protein, and decreased hepatic TG secretion (71).

Ren et al. compared mice fed high fat diet or high fructose diets. Liver lipid levels increased within 3 days and indicators of impaired glucose tolerance and insulin signaling were exhibited within one week in both groups. As expected, DNL and lipogenic gene expression were increased in fructose-fed mice and decreased in fat-fed mice. Interestingly, fructose feeding activated two endoplasmic reticulum stress pathways, but high fat feeding did not. The authors suggest that endoplasmic reticulum stress is involved in DNL *per se* rather than resulting from hepatic steatosis or insulin resistance (72).

Extensive recent work by Richard Johnson and co-workers suggests that fructose-induced increases of uric acid may contribute to the adverse effects of fructose. They report that in fructose-exposed human hepatocytes, uric acid upregulates fructokinase expression (73) and inhibits AMP-activated kinase activity (74), thus amplifying the lipogenic effects of fructose. A series of experiments involving hepatocytes, allopurinol-treated mice, and hyperuricemic patients with low BMI provides evidence that the lipogenic effects of fructose may be partially mediated through direct effects of uric acid to stimulate hepatic fat

accumulation (75). Most recently Tapia et al. studied 3 groups of rats receiving: 1. Uricase inhibitor treatment, 2. SSB, 3. Uricase inhibitor + SSB. Uricase inhibitor induced glomerular hypertension and SSB induced insulin resistance. In combination, they produced both effects, plus synergistic effects on systemic and glomerular pressure, plasma glucose, hepatic TG, and oxidative stress (76).

## Conclusion

The extent to which the adverse metabolic effects of dietary sugar consumption result from direct effects of fructose on lipid and carbohydrate metabolism, to indirect effects resulting from increased body weight and adiposity, or to direct metabolic actions that are exacerbated by weight gain, has not been determined.

However, the accumulating epidemiological evidence, direct clinical evidence, and the evidence suggesting plausible mechanisms support a role for sugar in the epidemics of metabolic syndrome, CVD and T2DM.

## Acknowledgments

The studies conducted by Dr. Havel's research group were supported with funding from NIH grants R01 HL-075675, 1R01 HL-091333, 1R01 HL-107256 and a Multi-campus Award from the University of California, Office of the President (UCOP #142691). These projects also received support from Grant Number UL1 RR024146 from the National Center for Research Resources (NCRR), a component of the National Institutes of Health (NIH), and NIH Roadmap for Medical Research. Kimber Stanhope is supported by a Building Interdisciplinary Research Careers in Women's Health award (K12 HD051958) funded by the National Institute of Child Health and Human Development (NICHD), Office of Research on Women's Health (ORWH), Office of Dietary Supplements (ODS), and the National Institute of Aging (NIA).

The authors thank James Graham, Marinelle Nuñez, Guoxia Chen, Vivien Lee, Hazel Lam, and Janet Peerson for their excellent ongoing technical support of the clinical studies, and Study Physicians, Drs. Steven Griffen, Andrew Bremer and Valentine Medici, as well as Nicole Mullen, R.N. and the nursing staff at CCRC for their dedicated nursing support.

## References

1. Johnson RK, Appel LJ, Brands M, Howard BV, Lefevre M, Lustig RH, et al. Dietary sugars intake and cardiovascular health: a scientific statement from the American Heart Association. *Circulation*. 2009; 120(11):1011–20. Epub 2009/08/26. [PubMed: 19704096]
2. DGAC. Report of the Dietary Guidelines Advisory Committee (DGAC) on the Dietary Guidelines for Americans, 2010. 2010. <http://www.cnpp.usdagov/DGAs2010-DGACReport.htm>
3. Rebollo A, Roglans N, Alegret M, Laguna JC. Way back for fructose and liver metabolism: Bench side to molecular insights. *World J Gastroenterol*. 2012; 18(45):6552–9. Epub 2012/12/14. [PubMed: 23236229]
4. Tappy L, Mittendorfer B. Fructose toxicity: is the science ready for public health actions? *Curr Opin Clin Nutr Metab Care*. 2012; 15(4):357–61. Epub 2012/05/24. [PubMed: 22617566]
5. Te Morenga L, Mallard S, Mann J. Dietary sugars and body weight: systematic review and meta-analyses of randomised controlled trials and cohort studies. *Bmj*. 2012; 346:e7492. Epub 2013/01/17. [PubMed: 23321486]
6. Klurfeld DM, Foreyt J, Angelopoulos TJ, Rippe JM. Lack of evidence for high fructose corn syrup as the cause of the obesity epidemic. *Int J Obes (Lond)*. 2012 Epub 2012/09/19.
- \*7. de Koning L, Malik VS, Kellogg MD, Rimm EB, Willett WC, Hu FB. Sweetened beverage consumption, incident coronary heart disease, and biomarkers of risk in men. *Circulation*. 2012; 125(14):1735–41. S1. Epub 2012/03/14. [PubMed: 22412070] In addition to reporting a positive association between consumption of SSB consumption and coronary heart disease and

intermediate biomarkers, this paper also reports that consumption of artificially sweetened beverage was not associated with any of these outcomes.

- \*8. Mattei J, Malik V, Hu FB, Campos H. Substituting homemade fruit juice for sugar-sweetened beverages is associated with lower odds of metabolic syndrome among Hispanic adults. *J Nutr.* 2012; 142(6):1081–7. Epub 2012/05/04. [PubMed: 22551801] This paper reports that substituting one serving of homemade fruit juice for instant SSB or regular soda was associated with ~30% lower odds of metabolic syndrome; a finding that warrants further investigation.
- \*9. Pan A, Malik VS, Schulze MB, Manson JE, Willett WC, Hu FB. Plain-water intake and risk of type 2 diabetes in young and middle-aged women. *Am J Clin Nutr.* 2012; 95(6):1454–60. Epub 2012/05/04. [PubMed: 22552035] In contrast to the finding from the study above, this study reported that consumption of fruit juice, as well as SSB, was associated with increased risk for T2DM.
10. Bhupathiraju SN, Pan A, Malik VS, Manson JE, Willett WC, van Dam RM, et al. Caffeinated and caffeine-free beverages and risk of type 2 diabetes. *Am J Clin Nutr.* 2013; 97(1):155–66. Epub 2012/11/16. [PubMed: 23151535]
- \*11. Siegel KR, Echouffo-Tcheugui JB, Ali MK, Mehta NK, Narayan KM, Chetty V. Societal correlates of diabetes prevalence: An analysis across 94 countries. *Diabetes Res Clin Pract.* 2012; 96(1):76–83. Epub 2011/12/23. [PubMed: 22189172] In addition to the association between SSB and increased incidence of diabetes, this paper also reported that each additional unit of fruit and vegetable availability was associated with 3% lower incidence of diabetes.
12. Wang JW, Mark S, Henderson M, O'Loughlin J, Tremblay A, Wortman J, et al. Adiposity and glucose intolerance exacerbate components of metabolic syndrome in children consuming sugar-sweetened beverages: QUALITY cohort study. *Pediatric obesity.* 2012 Epub 2012/11/23.
- \*13. Zgaga L, Theodoratou E, Kyle J, Farrington SM, Agakov F, Tenesa A, et al. The association of dietary intake of purine-rich vegetables, sugar-sweetened beverages and dairy with plasma urate, in a cross-sectional study. *PLoS ONE.* 2012; 7(6):e38123. Epub 2012/06/16. [PubMed: 22701608] Besides the positive association between SSB and uric acid, other findings of interest from this paper were that consumption of purine-rich vegetables were not associated with uric acid levels and that dairy consumption was negatively associated with uric acid levels.
- \*14. Lin WT, Huang HL, Huang MC, Chan TF, Ciou SY, Lee CY, et al. Effects on uric acid, body mass index and blood pressure in adolescents of consuming beverages sweetened with high-fructose corn syrup. *Int J Obes (Lond).* 2012 Epub 2012/08/15. This paper reports a positive association between SSB and uric acid, and also reports that the presence of obesity exacerbated the SSB-induced increases in uric acid. Recent work (reviewed in this paper) and a recent review from Richard Johnson and colleagues (Johnson, Seminars in nephrology, 2011) suggest a reverse of this relationship; that SSB-induced increases of uric acid may exacerbate obesity.
15. Pan A, Malik VS, Hao T, Willett WC, Mozaffarian D, Hu FB. Changes in water and beverage intake and long-term weight changes: results from three prospective cohort studies. *Int J Obes (Lond).* 2013 Epub 2013/01/16.
- \*16. Qi Q, Chu AY, Kang JH, Jensen MK, Curhan GC, Pasquale LR, et al. Sugar-sweetened beverages and genetic risk of obesity. *N Engl J Med.* 2012; 367(15):1387–96. Epub 2012/09/25. [PubMed: 22998338] The findings of this paper suggest that persons with genetic predisposition to obesity are especially vulnerable to the effects of SSB to promote body weight gain.
- \*17. Odegaard AO, Choh AC, Czerwinski SA, Towne B, Demerath EW. Sugar-sweetened and diet beverages in relation to visceral adipose tissue. *Obesity (Silver Spring).* 2012; 20(3):689–91. Epub 2011/09/09. [PubMed: 21901024] This paper provides support for our group's seminal finding that consumption of fructose can promote VAT accumulation (27).
- \*18. Pollock NK, Bundy V, Kanto W, Davis CL, Bernard PJ, Zhu H, et al. Greater fructose consumption is associated with cardiometabolic risk markers and visceral adiposity in adolescents. *J Nutr.* 2012; 142(2):251–7. Epub 2011/12/23. [PubMed: 22190023] This paper suggests that our finding that consumption of fructose promoted VAT accumulation in older, overweight/obese adults (27) is relevant to adolescents.
19. Sun SZ, Anderson GH, Flickinger BD, Williamson-Hughes PS, Empie MW. Fructose and nonfructose sugar intakes in the US population and their associations with indicators of metabolic syndrome. *Food and chemical toxicology : an international journal published for the British*

Industrial Biological Research Association. 2011; 49(11):2875–82. Epub 2011/09/06. [PubMed: 21889564]

- \*\*20. Maersk M, Belza A, Stodkilde-Jorgensen H, Ringgaard S, Chabanova E, Thomsen H, et al. Sucrose-sweetened beverages increase fat storage in the liver, muscle, and visceral fat depot: a 6-mo randomized intervention study. *Am J Clin Nutr*. 2012; 95(2):283–9. Epub 2011/12/30. [PubMed: 22205311] This paper reports the first direct experimental evidence that consuming excess sugar with an *ad libitum* diet increases liver TG content, as previous evidence came from an overfeeding study in which subjects were instructed to consume energy-balanced diets plus excess sugar (Le, *Am J Clin Nutr*, 2009). It also provides direct experimental evidence that corroborate our previous observation that fructose consumption can increase visceral adiposity (27).
- \*\*21. Aeberli I, Gerber PA, Hochuli M, Kohler S, Haile SR, Gouni-Berthold I, et al. Low to moderate sugar-sweetened beverage consumption impairs glucose and lipid metabolism and promotes inflammation in healthy young men: a randomized controlled trial. *Am J Clin Nutr*. 2011 Epub 2011/06/17. This paper is important for demonstrating that a moderate dose of sucrose promotes a more adverse lipid profile in as little as 3 weeks.
- \*\*22. Aeberli I, Hochuli M, Gerber PA, Sze L, Murer SB, Tappy L, et al. Moderate Amounts of Fructose Consumption Impair Insulin Sensitivity in Healthy Young Men: A randomized controlled trial. *Diabetes Care*. 2012 Epub 2012/08/31. The results in this paper are important in that they demonstrate a moderate dose of fructose affects hepatic insulin sensitivity in as little as 3 weeks, and they also corroborate our previous suggestion that fructose induces hepatic insulin resistance prior to whole body insulin resistance (Stanhope, *Curr Opin Lipidol*, 2008).
- \*\*23. Stanhope KL, Bremer AA, Medici V, Nakajima K, Ito Y, Nakano T, et al. Consumption of fructose and high fructose corn syrup increase postprandial triglycerides, LDL-cholesterol, and apolipoprotein-B in young men and women. *J Clin Endocrinol Metab*. 2011; 96(10):E1596–605. Epub 2011/08/19. [PubMed: 21849529] This is the first report of the effects of consumption of HFCS sweetened beverages (>24 h) on metabolic risk factors. Lipid risk factors for CVD (LDL, ApoB and sdLDL) were elevated within 2 weeks of consuming fructose- or HFCS-sweetened beverages in young (18–40 year old) men and women. To control for variations in diet and physical activity, the subjects in this study resided at the clinical research center for 48 hour prior to the baseline and 2-week intervention blood collections that were analyzed for lipid outcomes.
24. Silbernagel G, Machann J, Unmuth S, Schick F, Stefan N, Haring HU, et al. Effects of 4-week very-high-fructose/glucose diets on insulin sensitivity, visceral fat and intrahepatic lipids: an exploratory trial. *Br J Nutr*. 2011:1–8. Epub 2011/03/15.
25. Rao SS, Attaluri A, Anderson L, Stumbo P. Ability of the normal human small intestine to absorb fructose: evaluation by breath testing. *Clin Gastroenterol Hepatol*. 2007; 5(8):959–63. Epub 2007/07/13. [PubMed: 17625977]
26. Sevastianova K, Santos A, Kotronen A, Hakkarainen A, Makkonen J, Silander K, et al. Effect of short-term carbohydrate overfeeding and long-term weight loss on liver fat in overweight humans. *Am J Clin Nutr*. 2012; 96(4):727–34. Epub 2012/09/07. [PubMed: 22952180]
27. Stanhope KL, Schwarz JM, Keim NL, Griffen SC, Bremer AA, Graham JL, et al. Consuming fructose-sweetened, not glucose-sweetened, beverages increases visceral adiposity and lipids and decreases insulin sensitivity in overweight/obese humans. *J Clin Invest*. 2009; 119(5):1322–34. Epub 2009/04/22. [PubMed: 19381015]
28. Sun SZ, Empie MW. Fructose metabolism in humans - what isotopic tracer studies tell us. *Nutr Metab (Lond)*. 2012; 9(1):89. Epub 2012/10/04. [PubMed: 23031075]
29. Donnelly KL, Smith CI, Schwarzenberg SJ, Jessurun J, Boldt MD, Parks EJ. Sources of fatty acids stored in liver and secreted via lipoproteins in patients with nonalcoholic fatty liver disease. *J Clin Invest*. 2005; 115(5):1343–51. [PubMed: 15864352]
30. Vedala A, Wang W, Neese RA, Christiansen MP, Hellerstein MK. Delayed secretory pathway contributions to VLDL-triglycerides from plasma NEFA, diet, and de novo lipogenesis in humans. *J Lipid Res*. 2006; 47(11):2562–74. [PubMed: 16929033]
31. Hudgins LC, Parker TS, Levine DM, Hellerstein MK. A dual sugar challenge test for lipogenic sensitivity to dietary fructose. *J Clin Endocrinol Metab*. 2011; 96(3):861–8. Epub 2011/01/22. [PubMed: 21252253]

32. Parks EJ, Skokan LE, Timlin MT, Dingfelder CS. Dietary sugars stimulate fatty acid synthesis in adults. *J Nutr.* 2008; 138(6):1039–46. Epub 2008/05/22. [PubMed: 18492831]
- \*\*33. Cox CL, Stanhope KL, Schwarz JM, Graham JL, Hatcher B, Griffen SC, et al. Consumption of fructose-sweetened beverages for 10 weeks reduces net fat oxidation and energy expenditure in overweight/obese men and women. *Eur J Clin Nutr.* 2012; 66(2):201–8. Epub 2011/09/29. [PubMed: 21952692] The results of this paper suggest that fructose may promote liver lipid accumulation both directly, via DNL, and indirectly, via DNL-induced inhibition of fatty acid oxidation.
34. Samuel VT. Fructose induced lipogenesis: from sugar to fat to insulin resistance. *Trends Endocrinol Metab.* 2011; 22(2):60–5. Epub 2010/11/12. [PubMed: 21067942]
35. Bortolotti M, Dubuis J, Schneiter P, Tappy L. Effects of dietary protein on lipid metabolism in high fructose fed humans. *Clin Nutr.* 2012; 31(2):238–45. Epub 2011/10/25. [PubMed: 22019201]
- \*36. Cox CL, Stanhope KL, Schwarz JM, Graham JL, Hatcher B, Griffen SC, et al. Consumption of fructose- but not glucose-sweetened beverages for 10 weeks increases circulating concentrations of uric acid, retinol binding protein- 4, and gamma-glutamyl transferase activity in overweight/ obese humans. *Nutr Metab (Lond).* 2012; 9(1):68. Epub 2012/07/26. [PubMed: 22828276] This is the first study to measure and compare the 24-h uric acid profile following sustained consumption of fructose and glucose. The finding suggests that changes of 24-h uric acid exposure, which were markedly increased by only fructose, may be a more sensitive indicator of increased metabolic dysfunction than changes of fasting levels, which were increased by both fructose and glucose. The liver enzyme, gamma glutamyl transferase, and retinol binding protein-4 were also increased during fructose consumption.
37. Bombardier AS, Derebail VK, Shoham DA, Anderson CA, Steffen LM, Rosamond WD, et al. Sugar-sweetened soda consumption, hyperuricemia, and kidney disease. *Kidney international.* 2010; 77(7):609–16. Epub 2009/12/25. [PubMed: 20032963]
38. Chang WC. Dietary intake and the risk of hyperuricemia, gout and chronic kidney disease in elderly Taiwanese men. *The aging male : the official journal of the International Society for the Study of the Aging Male.* 2011; 14(3):195–202. Epub 2010/09/21.
39. Choi JW, Ford ES, Gao X, Choi HK. Sugar-sweetened soft drinks, diet soft drinks, and serum uric acid level: the Third National Health and Nutrition Examination Survey. *Arthritis and rheumatism.* 2008; 59(1):109–16. Epub 2008/01/01. [PubMed: 18163396]
- \*40. Cox CL, Stanhope KL, Schwarz JM, Graham JL, Hatcher B, Griffen SC, et al. Circulating concentrations of monocyte chemoattractant protein-1, plasminogen activator inhibitor-1, and soluble leukocyte adhesion molecule-1 in overweight/obese men and women consuming fructose- or glucose-sweetened beverages for 10 weeks. *J Clin Endocrinol Metab.* 2011; 96(12):E2034–8. Epub 2011/10/01. [PubMed: 21956423] The results from this study suggest the possibility that prolonged consumption of fructose may contribute to the development of metabolic syndrome via induction of specific proinflammatory and prothrombotic mediators.
- \*\*41. Stanhope KL, Griffen SC, Bremer AA, Vink RG, Schaefer EJ, Nakajima K, et al. Metabolic responses to prolonged consumption of glucose- and fructose-sweetened beverages are not associated with postprandial or 24-h glucose and insulin excursions. *Am J Clin Nutr.* 2011; 94(1):112–9. Epub 2011/05/27. [PubMed: 21613559] This study provides clear evidence that the adverse metabolic effects of fructose compared with glucose consumption are not explained by the glycemic index/glycemic load of the diets, nor mediated through postprandial glucose or insulin exposure.
42. Esfahani A, Wong JM, Mirrahimi A, Srichaikul K, Jenkins DJ, Kendall CW. The glycemic index: physiological significance. *J Am Coll Nutr.* 2009; 28(Suppl):439S–45S. Epub 2010/03/27. [PubMed: 20234030]
43. Brand-Miller J, Buyken AE. The glycemic index issue. *Curr Opin Lipidol.* 2012; 23(1):62–7. Epub 2011/12/14. [PubMed: 22157060]
- \*44. Jin R, Le NA, Liu S, Farkas Epperson M, Ziegler TR, Welsh JA, et al. Children with NAFLD are more sensitive to the adverse metabolic effects of fructose beverages than children without NAFLD. *J Clin Endocrinol Metab.* 2012; 97(7):E1088–98. Epub 2012/05/01. [PubMed: 22544914] The results from this study suggest that children with NAFLD may be more sensitive to the lipogenic effects of fructose consumption than children without NAFLD.



45. Teff KL, Elliott SS, Tschop M, Kieffer TJ, Rader D, Heiman M, et al. Dietary fructose reduces circulating insulin and leptin, attenuates postprandial suppression of ghrelin, and increases triglycerides in women. *J Clin Endocrinol Metab.* 2004; 89(6):2963–72. [PubMed: 15181085]
46. Teff KL, Grudziak J, Townsend RR, Dunn TN, Grant RW, Adams SH, et al. Endocrine and metabolic effects of consuming fructose- and glucose-sweetened beverages with meals in obese men and women: influence of insulin resistance on plasma triglyceride responses. *J Clin Endocrinol Metab.* 2009; 94(5):1562–9. Epub 2009/02/12. [PubMed: 19208729]
- \*47. Hasson RE, Adam TC, Davis JN, Kelly LA, Ventura EE, Byrd-Williams CE, et al. Randomized controlled trial to improve adiposity, inflammation, and insulin resistance in obese African-American and Latino youth. *Obesity (Silver Spring).* 2012; 20(4):811–8. Epub 2011/02/05. [PubMed: 21293446] This study demonstrated that reducing added sugar consumption improved insulin sensitivity in teenagers despite the absence of body weight or fat loss.
48. Goran MI, Walker R, Allayee H. Genetic-related and carbohydrate-related factors affecting liver fat accumulation. *Curr Opin Clin Nutr Metab Care.* 2012; 15(4):392–6. Epub 2012/05/24. [PubMed: 22617559]
49. Davis JN, Le KA, Walker RW, Vikman S, Spruijt-Metz D, Weigensberg MJ, et al. Increased hepatic fat in overweight Hispanic youth influenced by interaction between genetic variation in PNPLA3 and high dietary carbohydrate and sugar consumption. *Am J Clin Nutr.* 2010; 92(6):1522–7. Epub 2010/10/22. [PubMed: 20962157]
50. Sevastianova K, Kotronen A, Gastaldelli A, Perttila J, Hakkarainen A, Lundbom J, et al. Genetic variation in PNPLA3 (adiponutrin) confers sensitivity to weight loss-induced decrease in liver fat in humans. *Am J Clin Nutr.* 2011; 94(1):104–11. Epub 2011/04/29. [PubMed: 21525193]
51. Brymora A, Flisinski M, Johnson RJ, Goszka G, Stefanska A, Manitius J. Low-fructose diet lowers blood pressure and inflammation in patients with chronic kidney disease. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association.* 2012; 27(2):608–12. Epub 2011/05/27.
52. Maier IB, Stricker L, Ozel Y, Wagnerberger S, Bischoff SC, Bergheim I. A low fructose diet in the treatment of pediatric obesity: a pilot study. *Pediatrics international : official journal of the Japan Pediatric Society.* 2011; 53(3):303–8. Epub 2010/09/14. [PubMed: 20831653]
- \*\*53. de Ruyter JC, Olthof MR, Seidell JC, Katan MB. A trial of sugar-free or sugar-sweetened beverages and body weight in children. *N Engl J Med.* 2012; 367(15):1397–406. Epub 2012/09/25. [PubMed: 22998340] This blinded and randomized 18-month intervention comparing the effects of consuming one daily serving of SSB containing 104 kcals with consuming one serving of non-caloric sweetened beverage in 641 children provides strong evidence that SSB promotes weight gain in children. It also suggests that strategies that target reduction of SSB consumption in children may be effective in lowering the prevalence of pediatric overweight and obesity.
- \*54. Ebbeling CB, Feldman HA, Chomitz VR, Antonelli TA, Gortmaker SL, Osganian SK, et al. A randomized trial of sugar-sweetened beverages and adolescent body weight. *N Engl J Med.* 2012; 367(15):1407–16. Epub 2012/09/25. [PubMed: 22998339] This study also suggests that strategies that are effective for reducing SSB consumption may lower the prevalence of overweight/obesity in children.
55. Raben A, Vasilaras TH, Moller AC, Astrup A. Sucrose compared with artificial sweeteners: different effects on ad libitum food intake and body weight after 10 wk of supplementation in overweight subjects. *Am J Clin Nutr.* 2002; 76(4):721–9. Epub 2002/09/27. [PubMed: 12324283]
56. Reid M, Hammersley R, Hill AJ, Skidmore P. Long-term dietary compensation for added sugar: effects of supplementary sucrose drinks over a 4-week period. *Br J Nutr.* 2007; 97(1):193–203. Epub 2007/01/16. [PubMed: 17217576]
57. Tordoff MG, Alleva AM. Effect of drinking soda sweetened with aspartame or high-fructose corn syrup on food intake and body weight. *Am J Clin Nutr.* 1990; 51(6):963–9. Epub 1990/06/01. [PubMed: 2349932]
- \*58. Ng J, Stice E, Yokum S, Bohon C. An fMRI study of obesity, food reward, and perceived caloric density. Does a low-fat label make food less appealing?. *Appetite.* 2011; 57(1):65–72. Epub 2011/04/19. Using fMRI, the investigators demonstrated that areas of the brain associated with reward valuation were more activated when subjects consumed a milk shake labeled “regular”

compared with an identical milk shake labeled “low-fat”. This interesting finding suggests that preconceptions regarding diet composition, even when they are inaccurate, can affect the brain responses to the diet, and therefore possibly energy intake as well.

- \*\*59. Page KA, Chan O, Arora J, Belfort-Deaguiar R, Dzuira J, Roehmholdt B, et al. Effects of fructose vs glucose on regional cerebral blood flow in brain regions involved with appetite and reward pathways. *JAMA*. 2013; 309(1):63–70. Epub 2013/01/03. [PubMed: 23280226] This study shows that glucose compared with fructose ingestion results in differential brain activation patterns, which may be involved in the effects of SSB to promote body weight gain.
60. Purnell JQ, Klopfenstein BA, Stevens AA, Havel PJ, Adams SH, Dunn TN, et al. Brain functional magnetic resonance imaging response to glucose and fructose infusions in humans. *Diabetes Obes Metab*. 2011; 13(3):229–34. Epub 2011/01/06. [PubMed: 21205113]
61. Berkey CS, Rockett HR, Field AE, Gillman MW, Colditz GA. Sugar-added beverages and adolescent weight change. *Obes Res*. 2004; 12(5):778–88. Epub 2004/05/29. [PubMed: 15166298]
62. Ludwig DS, Peterson KE, Gortmaker SL. Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet*. 2001; 357(9255):505–8. Epub 2001/03/07. [PubMed: 11229668]
63. Mozaffarian D, Hao T, Rimm EB, Willett WC, Hu FB. Changes in diet and lifestyle and long-term weight gain in women and men. *N Engl J Med*. 2011; 364(25):2392–404. Epub 2011/06/24. [PubMed: 21696306]
64. Hallfrisch J, Ellwood KC, Michaelis OEt, Reiser S, O’Dorisio TM, Prather ES. Effects of dietary fructose on plasma glucose and hormone responses in normal and hyperinsulinemic men. *J Nutr*. 1983; 113(9):1819–26. [PubMed: 6350544]
65. Reiser S, Bickard MC, Hallfrisch J, Michaelis OEt, Prather ES. Blood lipids and their distribution in lipoproteins in hyperinsulinemic subjects fed three different levels of sucrose. *J Nutr*. 1981; 111(6):1045–57. [PubMed: 6940954]
66. Reiser S, Bohn E, Hallfrisch J, Michaelis OEt, Keeney M, Prather ES. Serum insulin and glucose in hyperinsulinemic subjects fed three different levels of sucrose. *Am J Clin Nutr*. 1981; 34(11):2348–58. [PubMed: 7030048]
67. Reiser S, Hallfrisch J, Michaelis OEt, Lazar FL, Martin RE, Prather ES. Isocaloric exchange of dietary starch and sucrose in humans. I. Effects on levels of fasting blood lipids. *Am J Clin Nutr*. 1979; 32(8):1659–69.
68. Reiser S, Handler HB, Gardner LB, Hallfrisch JG, Michaelis OEt, Prather ES. Isocaloric exchange of dietary starch and sucrose in humans. II. Effect on fasting blood insulin, glucose, and glucagon and on insulin and glucose response to a sucrose load. *Am J Clin Nutr*. 1979; 32(11):2206–16.
69. Reiser S, Powell AS, Scholfield DJ, Panda P, Fields M, Canary JJ. Day-long glucose, insulin, and fructose responses of hyperinsulinemic and nonhyperinsulinemic men adapted to diets containing either fructose or high-amylose cornstarch. *Am J Clin Nutr*. 1989; 50(5):1008–14. [PubMed: 2683715]
- \*\*70. Ishimoto T, Lanaspas MA, Le MT, Garcia GE, Diggle CP, Maclean PS, et al. Opposing effects of fructokinase C and A isoforms on fructose-induced metabolic syndrome in mice. *Proc Natl Acad Sci U S A*. 2012; 109(11):4320–5. Epub 2012/03/01. [PubMed: 22371574] This study demonstrates that the metabolism of fructose through fructokinase C is necessary to generate adverse metabolic effects. It also demonstrates that when fructose flux through fructokinase C is increased due to knockout of fructokinase A, adverse metabolic effects of fructose are exacerbated.
- \*71. Erion DM, Popov V, Hsiao JJ, Vatner D, Mitchell K, Yonemitsu S, et al. The role of the carbohydrate response element-binding protein in male fructose-fed rats. *Endocrinology*. 2013; 154(1):36–44. Epub 2012/11/20. [PubMed: 23161873] This study suggests that ChREBP is involved in activation of lipogenesis in fructose-fed rats, and that it also has a role in promoting hepatic TG secretion.
- \*72. Ren LP, Chan SM, Zeng XY, Laybutt DR, Iseli TJ, Sun RQ, et al. Differing endoplasmic reticulum stress response to excess lipogenesis versus lipid oversupply in relation to hepatic steatosis and insulin resistance. *PLoS ONE*. 2012; 7(2):e30816. Epub 2012/02/23. [PubMed: 22355328] The results of this study suggests that endoplasmic reticulum stress is associated with upregulation of DNL, rather than resulting from hepatic steatosis or insulin resistance. Thus,




development of fatty liver due to fructose-induced DNL may be more problematic than fatty liver induced by high fat diet.

73. Lanaspá MA, Sanchez-Lozada LG, Cicerchi C, Li N, Roncal-Jimenez CA, Ishimoto T, et al. Uric acid stimulates fructokinase and accelerates fructose metabolism in the development of fatty liver. PLoS ONE. 2012; 7(10):e47948. Epub 2012/11/01. [PubMed: 23112875]
74. Lanaspá MA, Cicerchi C, Garcia G, Li N, Roncal-Jimenez CA, Rivard CJ, et al. Counteracting roles of AMP deaminase and AMP kinase in the development of fatty liver. PLoS ONE. 2012; 7(11):e48801. Epub 2012/11/16. [PubMed: 23152807]
- \*75. Lanaspá MA, Sanchez-Lozada LG, Choi YJ, Cicerchi C, Kanbay M, Roncal-Jimenez CA, et al. Uric acid induces hepatic steatosis by generation of mitochondrial oxidative stress: potential role in fructose-dependent and -independent fatty liver. J Biol Chem. 2012; 287(48):40732–44. Epub 2012/10/05. [PubMed: 23035112] This study suggests that the lipogenic effects of fructose are exacerbated by high uric acid levels.
- \*76. Tapia E, Cristobal M, Garcia-Arroyo FE, Soto V, Monroy-Sanchez F, Pacheco U, et al. Synergistic Effect of Uricase Blockade Plus Physiologic Amounts of Fructose-Glucose on Glomerular Hypertension and Oxidative Stress in Rats. Am J Physiol Renal Physiol. 2013 Epub 2013/01/11. This study provides evidence that SSB and high uric acid levels interact to produce more adverse effects in rats than either SSB or high uric acid levels alone. While these are important mechanistic results, they have potential clinical implications for hyperuricemic individuals.


### Key points

- Recently published epidemiological studies that provide evidence that sugar consumption is associated with metabolic disease.
- Three recent clinical studies, which investigated the effects of consuming relevant doses of sucrose or high fructose corn syrup along with *ad libitum* diets, provide evidence that consumption of these sugars increase risk factors for cardiovascular disease and metabolic syndrome.
- Mechanistic studies suggest that the adverse effects of sugar consumption result from the rapid hepatic metabolism of fructose catalyzed by fructokinase C, which generates substrate for *de novo* lipogenesis and leads to increased uric acid levels.
- Recent clinical studies investigating the effects of consuming less sugar, via educational interventions or by substitution of sugar-sweetened beverages for non-calorically sweetened beverages, provide evidence that such strategies have beneficial effects on risk factors for metabolic disease or on BMI in children.



**Figure 1. Effect of body weight gain on the changes of fasting cholesterol, LDL and apoB in subjects consuming sugar**

Changes in fasting lipid outcomes in subjects who gained and did not gain body weight while consuming 25%E HFCS-, fructose-, or glucose-sweetened beverages for 2 weeks with ad libitum diets (\* $P < 0.05$ , \*\* $P < 0.01$ ; 2 week vs baseline).



**Figure 2. Two mechanisms by which sugar increases metabolic risk**

Consumption of sugar increases risk for metabolic disease via the direct effects of fructose on lipid and carbohydrate metabolism. Consumption of sugar may also promote body weight gain. The increased body weight/adiposity further increases risk for metabolic disease. Thus consumption of sugar increases risk for metabolic disease both directly and via effects to promote weight gain.